

Role of protected areas in climate change mitigation and biodiversity conservation

Technical Brief | UNFCCC COP 28
November 2023

Prepared by the World Commission on Protected Areas (WCPA) of the International Union for the Conservation of Nature (IUCN)



Key Messages

1. Widespread consensus has emerged that biodiversity loss and climate change are twin crises that must be addressed together to achieve success in either one.
2. Protected and Conserved Areas (PAs) are the most effective tool to address both biodiversity loss and climate change within a time-frame that reflects the required urgency.
3. New PAs, the expansion of existing PAs and support for OECMs can target places where carbon richness and high biodiversity overlap to create 'carbon stabilization' areas.

Broad Policy Consensus

There is general consensus that biodiversity loss and climate change are twin crises requiring integrated, comprehensive and holistic approaches. As aptly articulated by António Guterres, Secretary-General of the United Nations, the twin global challenges of climate change and biodiversity loss have to be tackled in a more coordinated manner: "climate change threatens to undermine all efforts to conserve and sustainably manage biodiversity and [that] nature itself offers some of the most effective solutions to avert the worst impacts of a warming planet". The creation of new Protected areas (PAs), the expansion of existing ones, protecting and establishing conservation corridors that connect PAs, and better PA management are the first and most effective policy tools to address these global crises.

UNFCCC COP 27 underlined "...the urgent need to address in a comprehensive and synergistic manner, the interlinked global crises of climate change and biodiversity loss in the broader context of achieving the Sustainable Development Goals, as well as the vital importance of protecting, conserving, restoring and sustainably using nature and ecosystems for effective and sustainable climate action" (Dec.1/CMA 4, para 1).

Likewise, the United Nations Convention on Biological Diversity (CBD) recognized the importance of the climate change/biodiversity nexus [3], including in the [Kunming Declaration](#) from COP15 Part 1 and the Kunming-Montreal Global Biodiversity Framework (target 8) [4]. A nature-positive world, as promoted by the 95 country signatories to [the Leaders Pledge for Nature](#), embodies this collective understanding of the interdependence of biodiversity loss, ecosystem integrity, climate change impacts and human well-being [5].

Status of Biodiversity and Climate Change Targets

Some significant successes have been achieved in protecting natural ecosystems and biodiversity. From 2010 to 2023 terrestrial protected areas have increased from about 10% of the land and inland waters to over 16.05%, and marine protected areas have increased from 3% to 8.17%. If Other Effective Conservation Measures (OECMs) are included, the total terrestrial and inland waters protected stands at 17.23% and marine protected areas at 8.28% [6].



Example of a carbon-rich, high biodiversity tropical rainforest Yasuni National Park, Ecuador Photo: Maris Maskalanslistock

Forty-two percent of the area now within PAs has been added since 2010. In spite of these successes, most indicators of ecosystems and biodiversity are showing rapid decline, with 75% of the land surface significantly altered, 66% of the ocean area experiencing increasing cumulative impacts, over 85% of wetland area having been lost, and 25% of assessed animal and plant species threatened [7]. As well, the tropics lost 10% more primary rainforest in 2022 than in 2021, the year after the Glasgow Leaders' Declaration on Forests and Land Use to halt and reverse forest loss by the end of the decade [8]. Climate change is now one of the top five drivers of biodiversity loss globally [7].

Since the Paris Agreement in 2015, greenhouse gas (GHG) emissions increased every year, except for a small decline from 2019 to 2020 due to the COVID-19 pandemic.

As well, the concentration of CO₂ in the atmosphere continued to grow [9] and atmospheric concentrations of CO₂ reached an all-time high in September 2022 of 419 parts per million (ppm), from 280 ppm in the mid-1700s. Natural carbon sinks on land and in the ocean absorbed about half of the CO₂ emitted each year between 2011 and 2020. The other half accumulated in the atmosphere [10].

All pathways to attaining the Paris Agreement target of limiting average global temperature increases to well under 2°C above pre-industrial levels require protecting the global carbon sinks and reservoirs in natural ecosystems, in addition to rapid and drastic declines in GHG emissions. One fundamental and well-recognized tool is the protection of carbon-rich ecosystems such as primary forests, peatlands, grasslands, coastal blue carbon and marine biota, as well as the restoration of lost and damaged terrestrial, freshwater, coastal and marine ecosystems. These ecosystems are often referred to as 'carbon-rich' because they sequester and store more carbon than other ecosystems.

Supporting Evidence

The science agencies supporting the UN Rio Conventions all identified the important role of protected areas in realizing the synergies between the two conventions. The [sixth assessment report \(AR6\)](#) of the Intergovernmental Panel on Climate Change (IPCC) acknowledged area-based conservation, protection and restoration of ecosystems as effective strategies to reduce the vulnerability of biodiversity and humans to climate change [11]. It also concluded that actions to protect ecosystems with a high biodiversity value, such as forests, and particularly primary forests, coastal vegetated wetlands, peatlands, savannas and grasslands offer the highest total and per area mitigation value of any action in the Agriculture, Forestry and Other Land Uses (AFOLU) sector. And at the lowest cost.

The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystems Services (IPBES), in its Global Assessment Report recognizes the increasing impacts of climate change on biodiversity and the unlikelihood of reaching the Sustainable Development Goals or the Paris Agreement without addressing the relationship between nature and people. A joint IPBES-IPCC workshop on Biodiversity and Climate Change has also demonstrated the interdependence of biodiversity and climate change and the need to explore solutions that address both crises [12]. Recent analyses, drawing on multiple lines of evidence, suggest that maintaining the resilience of biodiversity and ecosystem services at a global scale depends on effective and equitable conservation of approximately 30 to 50% of Earth's land, freshwater and oceans [13].

Role of Protected Areas

The purpose of protected areas has evolved over time. The original concept of protecting scenic landscapes, culturally important areas and areas where people can connect to nature, was supplanted by the concept of maintaining populations of iconic and threatened species. As threats from climate change have intensified, there has been an increasing understanding of the value of PAs in protecting carbon sinks and stores, as well as other ecosystem services.

[The Protected and Conserved Areas Joint Statement on Climate Change and Biodiversity Crises](#), signed at the UNFCCC COP26, recognizes this expanded role for PAs. It encourages "better conservation and restoration of nature, especially of carbon-rich ecosystems, and better connectivity of these spaces to become the heart and anchors of nature recovery networks globally" [14].

Protecting existing carbon-rich ecosystems, such as primary forests, peatlands, grasslands, mangroves, seagrass beds and salt marshes has multiple benefits for biodiversity and people. Emissions benefits are achieved quickly (i.e. within a time-frame compatible with the urgency of the climate and biodiversity crises) and the cost per tonne of CO₂ is low [15]. Much of the carbon in these ecosystems has accumulated over centuries or longer: if released by human activities it is essentially irreplaceable in a time-frame that can meet the urgent need to reduce greenhouse gas emissions. About 23% of this irrecoverable carbon is found within PAs, with about half of it concentrated in just 3.3% of the planet's lands [16,17]. Globally, forests contain the most stored carbon, partly because of their widespread extent. Mangroves, salt marshes and seagrass beds contain more stored carbon on an area basis than most forests, although their limited geographic extent makes them less important on a global scale (Figure 1).





Figure 1: Estimates of global carbon storage in selected ecosystems. Sources for estimates are: Coastal blue carbon (mangroves, salt marshes and seagrass beds) [18] (Macreadie et al 2021); Grasslands [19] (Lorenz and Lal 2018); Peatlands [20] (Strack et al. 2022)

While areas rich in biodiversity and high in carbon do not always overlap, it is always the case that carbon retention in ecosystems is dependent of ecological integrity, which in turn is underpinned by an ecosystem's natural biodiversity. In both cases, protection offers high mitigation and biodiversity value. Where they do overlap, there is potential to significantly improve the protection of species rich ecosystems. In one global analysis, Soto-Navarro et al (2020) [22] found an overlap of 38% between carbon richness and a biodiversity index that prioritizes areas of high biodiversity (measured as high species richness, range-size rarity), high local intactness and high habitat condition, with 12% of the overlap area falling within existing PAs.

The same authors found only a 5% overlap between carbon richness and a biodiversity index that prioritizes high local biodiversity, low average habitat condition and high threats to biodiversity, but 21% of the overlap area fell within existing PAs (Figure 2). There is potential for increasing protected areas where the overlap is high and protection is low.

Protected areas have been estimated to store about 15.2% of terrestrial carbon stocks and to sequester about 20% of the carbon sequestered by all land ecosystems [23]. A similar analysis has not been made for Marine Protected Areas (MPAs), although they are believed to be significant, considering that the oceans have absorbed 20-25% of atmospheric carbon dioxide since 2008 [24].

Figure 2. The top map depicts the area of overlap between areas of high local biodiversity (high species richness, range- size rarity), high local intactness and high average habitat condition with carbon-richness. The bottom map depicts the area of overlap between areas of high local biodiversity, low average habitat condition and high threats with carbon richness. The dark brown areas depict the areas of highest overlap. Reprinted with permission [22].



Protected and Conserved Areas, as well as the conservation corridors that connect them [20]: i) provide the ecosystem integrity on which the long-term persistence of ecosystems depends; ii) enhance the resilience of natural ecosystems to change, and especially the disturbances resulting from climate change; and iii) reduce the vulnerability of ecosystems to multiple pressures across a landscape, including providing a buffer against global tipping points. They are thus a critically important strategy for retaining carbon in biosphere carbon reservoirs and supporting resilient, long lived carbon recovery in natural ecosystems.

Nationally Determined Contributions (NDCs) to the Paris Agreement and Protected Areas

In the first iteration of Nationally Determined Contributions (NDCs) to the Paris Agreement, 67 countries (37% of the NDCs examined) explicitly mentioned terrestrial protected areas or other conserved areas as part of their contribution to reduce greenhouse gas emissions, although few offered specific numbers on how much GHG reduction would be achieved by PAs [27]. A similar analysis on the updated NDCs remains to be done, but the role of nature in GHG emissions is receiving increasing prominence in discussions at UNFCCC, so we can presume that PAs will also be more prominent in updated NDCs.

Unlike in the terrestrial realm, few countries included MPAs in their NDCs to the Paris Agreement, although the potential for MPAs to provide both mitigation and adaptation benefits is well-documented. Oceans play a significant role in buffering climate change - by absorbing more than 90 percent of the temperature increases caused by global warming and 50 percent of all the carbon emissions since the Industrial Revolution [28,29].

However, this is not sustainable given the cost to marine ecosystems such as deoxygenation [30], ocean acidification and ocean temperature increases [31,32]. It has been estimated that between 0.02 to 0.65 billion tonnes (Gt) of CO₂e/year of GHG emissions could be sequestered and stored by increasing the protection and restoration of coastal ecosystems (mangroves, salt marshes and seagrasses) [31]. MPAs created and managed using climate information can enhance marine ecosystem resilience by:

- Enhancing connectivity between populations, species and ecosystems, including protecting species that migrate or travel over large distances;
- Providing climate refugia or safe havens for vulnerable species and ecosystems, and protecting populations of species important for food security;
- Improving the adaptive capacity of species, given the evidence that species and genetic diversity are often greater in marine reserves compared to harvested areas;
- Recognizing and supporting Other Effective Area-based Conservation Measures (OECMs) to protect key habitats, provide connectivity and enhance the resilience of coastal and marine ecosystem.

This IUCN WCPA Technical Note should be cited as:

Smith, R., Young, V. 2023. Role of Protected Areas in Climate Change Mitigation and Biodiversity Conservation. Technical Note SeriesNo. 8, Updated. Gland, Switzerland: IUCN WCPA. 6pp.

The views expressed in this publication do not necessarily reflect those of IUCN or other participating organisations.

References Cited

1. UNFCCC. Report of the Conference of the Parties on its twenty-fifth session, held in Madrid from 2-15 December 2019. Action taken by the Conference of the Parties at its twenty-fifth session. Chile Madrid Time for Action. Decisions 1/CP.25 (para 15). (UNFCCC, Madrid, Spain, 2019).
2. UNFCCC. Glasgow Climate Pact. Decisions 1/CMP.16. (2021).
3. Campbell, A. et al. Review of the Literature on the Links Between Biodiversity and Climate Change: Impacts, adaptation, and mitigation. Vol. Technical Series No. 42 124 (Secretariat of the Convention on Biological Diversity, 2009).
4. Convention on Biological Diversity. Kunming-Montreal Global Biodiversity Framework. (Montreal, Canada, 2022).
5. Leaders' Pledge 4 Nature. United to reverse biodiversity loss by 2030 for sustainable development. (New York, 2020).
6. WCMC. Protected Planet, <<https://www.protectedplanet.net/en>> (2023).
7. Secretariat of the Convention on Biological Diversity. Global Biodiversity Outlook 5. (Montreal, 2020).
8. Weisse, M., Goldman, E. & Carter, S. Forest Pulse: The Latest on the World's Forests. (2023).
9. UNEP. Emissions Gap Report: The Closing Window - Climate crisis calls for rapid transformation of societies. (Nairobi, Kenya, 2022).
10. NOAA. Climate Change: Atmospheric Carbon Dioxide. (2022).
11. Pörtner, H.-O. et al. WGII contribution to the IPCC Sixth Assessment Report (AR6), climate change 2022: Impacts, adaptation and vulnerability; Summary for policymakers. (Switzerland, 2022).
12. Nabuurs, G.-J. et al. in IPCC Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Vol. Chapter 7 (eds P.R. Shukla et al.) (Cambridge University Press, 2022).
13. Woodley, S. et al. A review of evidence for area-based conservation targets for the post-2020 global biodiversity network. PARKS 25, 31-46 (2019). <https://doi.org/https://doi.org/10.2305/IUCN.CH.2019.PARKS-25-2SW2.en>
14. Protected Areas Agencies. Protected and Conserved Areas Joint Statement on Climate Change and Biodiversity Crises. (UNFCCC, Glasgow, Scotland, 2021).
15. Cook-Patton, S. C. et al. Protect, manage and then restore lands for climate mitigation. Nature Climate Change 11, 1027-1034 (2021). <https://doi.org/https://doi.org/10.1038/s41558-021-01198-0>
16. Goldstein, A. et al. Protecting irrecoverable carbon in Earth's ecosystems. Nature Climate Change 10, 287-295 (2020). <https://doi.org/https://doi.org/10.1038/s41558-020-0738-8>
17. Noon, M. L. et al. Mapping the irrecoverable carbon in Earth's ecosystems. Nature Sustainability (2021). <https://doi.org/https://doi.org/10.1038/s41893-021-00803-6>
18. Macreadie, P. I. et al. Blue carbon as a natural climate solution. Nature Reviews Earth & Environment 2, 826-839 (2021). <https://doi.org/https://doi.org/10.1038/s43017-021-00224-1>
19. Lorenz, K. & Lal, R. in Carbon Sequestration in Agricultural Ecosystems(eds Klaus Lorenz & Rattan Lal)175-209 (Springer International Publishing, 2018).
20. Strack, M., Davidson, S. J., Hirano, T. & Dunn, C. The potential of peatlands as nature-based climate solutions. Current Climate Change Reports, 1-12 (2022). <https://doi.org/https://doi.org/10.1007/s40641-022-00183-9>
21. Woods Hole Climate Research Center et al. Policy Briefs: Primary Tropical, Temperate and Boreal Forests: Critical Stores of Carbon, Biodiversity and Freshwater. (2020).
22. Soto-Navarro, C. et al. Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action. Philosophical Transactions of the Royal Society B: Biological Sciences 375, 20190128 (2020). <https://doi.org/https://doi.org/10.1098/rstb.2019.0128>
23. Meilillo, J. M. et al. Protected areas' role in climate-change mitigation. Ambio 45, 133-145 (2016). <https://doi.org/https://doi.org/10.1007/s13280-015-0693-1>
24. Le Quéré, C. et al. Global Carbon Budget 2018. Earth System Science Data (Online) 10, 2141-2194 (2018). <https://doi.org/https://doi.org/10.5194/essd-10-2141-2018>
25. Graham, V. et al. Southeast Asian protected areas are effective in conserving forest cover and forest carbon stocks compared to unprotected areas. Sci Rep 11, 23760 (2021). <https://doi.org/https://doi.org/10.1038/s41598-021-03188-w>
26. Zhu, Z. et al. Conservation of carbon resources and values on public lands: A case study from the National Wildlife Refuge System. PloS one 17, e0262218 (2022). <https://doi.org/https://doi.org/10.1371/journal.pone.0262218>
27. Hehmeyer, A., Vogel, J., Martin, S. & Bartlett, R. Enhancing Nationally Determined Contributions Through Protected Areas. 1-44 (Washington DC, 2019).
28. NASA. Vital Signs of the Planet: Ocean Warming. (2022).
29. Sabine, C. L. et al. The oceanic sink for anthropogenic CO₂. science 305, 367-371 (2004).
30. Laffoley, D. & Baxter, J. M. Ocean deoxygenation: Everyone's problem-Causes, impacts, consequences and solutions. 1-580 (IUCN, 2019).
31. Northrop, E. et al. Enhancing Nationally Determined Contributions: Opportunities for Ocean-Based Climate Action. World Resources Institute: Washington, DC, USA (2021).
32. Laffoley, D. D. A. & Baxter, J. M. Explaining ocean warming: Causes, scale, effects and consequences. (IUCN Gland, Switzerland, 2016).